



ON PROJECTIVE φ -RECURRENT KENMOTSU MANIFOLDS

C.S. Bagewadi and Venkatesha*

ABSTRACT

In this paper we study a projective φ -recurrent Kenmotsu manifold and show that projective φ -recurrent Kenmotsu manifold having a non-zero constant sectional curvature is locally projective φ -symmetric.

MSC (2000): 53C05, 53C20, 53C25, 53D15.

Keywords and phrases: projective φ -recurrent, Kenmotsu manifold, sectional curvature, locally projective φ -symmetric.

1. Introduction

In 1977, T. Takahashi [5] introduced the notion of locally φ -symmetric Sasakian manifold and obtain few of its interesting properties. The authors [2] and [7], have extended this notion to 3-dimensional Kenmotsu and trans-Sasakian manifolds respectively. Also the authors [3] and [6] studied φ -recurrent Sasakian and Kenmotsu manifold respectively. In this paper we study a projective φ -recurrent Kenmotsu manifold and obtain some interesting results. Here we show that a projective φ -recurrent Kenmotsu manifold is an Einstein manifold and a projective φ -recurrent Kenmotsu manifold having a non-zero constant sectional curvature is locally projective φ -symmetric.

* Department of Mathematics and Computer Science, Kuvempu University, Jnana Sahyadri - 577 451, Shimoga, Karnataka, India

2. Preliminaries

Let $M^{2n+1}(\varphi, \xi, \eta, g)$ be a Kenmotsu manifold with the structure (φ, ξ, η, g) . Then the following relations hold [4]:

$$(2.1) \quad \varphi^2 = -I + \eta \otimes \xi$$

$$(2.2) \quad (a) \ \eta(\xi) = 1, \quad (b) \ g(X, \xi) = \eta(X), \quad (c) \ \varphi\xi = 0 \quad (d) \ \eta \circ \varphi = 0$$

$$(2.3) \quad g(\varphi X, \varphi Y) = g(X, Y) - \eta(X)\eta(Y),$$

$$(2.4) \quad (\nabla_X \varphi)Y = -g(X, \varphi Y)\xi - \eta(Y)\varphi X,$$

$$(2.5) \quad (a) \ \nabla_X \xi = X - \eta(X)\xi, \quad (b) \ (\nabla_X \eta)Y = g(X, Y) - \eta(X)\eta(Y)$$

$$(2.6) \quad R(X, Y)\xi = \eta(X)Y - \eta(Y)X$$

$$(2.7) \quad R(X, \xi)Y = g(X, Y)\xi - \eta(Y)X$$

$$(2.8) \quad \eta(R(X, Y)Z) = g(X, Z)\eta(Y) - g(Y, Z)\eta(X)$$

$$(2.9) \quad S(X, \xi) = -2n\eta(X)$$

$$(2.10) \quad S(\varphi X, \varphi Y) = S(X, Y) + 2n\eta(X)\eta(Y)$$

for all vector fields X, Y, Z , where ∇ denotes the operator of covariant differentiation with respect to g , φ is a $(1, 1)$ tensor field, S is the Ricci tensor of type $(0, 2)$ and R is the Riemannian curvature tensor of the manifold.

Definition 2.1. [2] A Kenmotsu manifold is said to be locally φ -symmetric if

$$\varphi^2((\nabla_W R)(X, Y)Z) = 0, \quad (2.11)$$

for all vector fields X, Y, Z, W orthogonal to ξ .

Definition 2.2. A Kenmotsu manifold is said to be locally projective φ -symmetric if

$$\varphi^2((\nabla_W P)(X, Y)Z) = 0, \quad (2.12)$$

for all vector fields X, Y, Z, W orthogonal to ξ .

Definition 2.3. A Kenmotsu manifold is said to be projective φ -recurrent manifold if there exists a non-zero 1-form A such that

$$\varphi^2((\nabla_w P)(X, Y)Z) = A(W)P(X, Y)Z \quad (2.13)$$

for arbitrary vector fields X, Y, Z, W , where P is a projective curvature tensor given by [1],

$$P(X, Y)Z = R(X, Y)Z - \frac{1}{2n}[S(Y, Z)X - S(X, Z)Y] \quad (2.14)$$

If the 1-form A vanishes, then the manifold reduces to a locally projective φ -symmetric manifold.

3. Projective φ -recurrent Kenmotsu Manifold

Let us consider a Projective φ -recurrent Kenmotsu manifold. Then by virtue of (2.1) and (2.13) we have

$$-(\nabla_w P)(X, Y)Z + \eta((\nabla_w P)(X, Y)Z)\xi = A(W)P(X, Y)Z \quad (3.1)$$

from which it follows that

$$-g((\nabla_w P)(X, Y)Z, U) + \eta((\nabla_w P)(X, Y)Z)\eta(U) = A(W)g(P(X, Y)Z, U) \quad (3.2)$$

Again from (3.1) we have

$$(\nabla_w P)(X, Y)Z = \eta((\nabla_w P)(X, Y)Z)\xi - A(W)P(X, Y)Z \quad (3.3)$$

This implies,

$$\begin{aligned} (\nabla_w R)(X, Y)Z &= \eta((\nabla_w R)(X, Y)Z)\xi - A(W)R(X, Y)Z \\ &+ \frac{1}{2n} [(\nabla_w S)(Y, Z)X - (\nabla_w S)(X, Z)Y] \\ &- \frac{1}{2n} [(\nabla_w S)(Y, Z)\eta(X) - (\nabla_w S)(X, Z)\eta(Y)]\xi \\ &+ \frac{1}{2n} A(W) [S(Y, Z)X - S(X, Z)Y] \end{aligned} \quad (3.4)$$

From (3.4) and the Bianchi identity we get

$$\begin{aligned}
& A(W)\eta(R(X, Y)Z) + A(X)\eta(R(Y, W)Z) + A(Y)\eta(R(W, X)Z) \\
&= \frac{1}{2n} A(W)[S(Y, Z)\eta(X) - S(X, Z)\eta(Y)] \\
&+ \frac{1}{2n} A(X)[S(W, Z)\eta(Y) - S(Y, Z)\eta(W)] \\
&+ \frac{1}{2n} A(Y)[S(X, Z)\eta(W) - S(W, Z)\eta(X)] \tag{3.5}
\end{aligned}$$

By virtue of (2.8) we obtain from (3.5) that

$$\begin{aligned}
& A(W)[\eta(Y)g(X, Z) - \eta(X)g(Y, Z) + A(X)[\eta(W)g(Y, Z) - \eta(Y)g(W, Z)] \\
&+ A(Y)[\eta(X)g(W, Z) - \eta(W)g(X, Z)] \\
&= \frac{1}{2n} A(W)[S(Y, Z)\eta(X) - S(X, Z)\eta(Y)] \\
&+ \frac{1}{2n} A(X)[S(W, Z)\eta(Y) - S(Y, Z)\eta(W)] \\
&+ \frac{1}{2n} A(Y)[S(X, Z)\eta(W) - S(W, Z)\eta(X)] \tag{3.6}
\end{aligned}$$

Putting $Y = Z = e_i$ in (3.6) and taking summation over i , $1 \leq i \leq 2n+1$, we get

$$A(W)\eta(X) = A(X)\eta(W), \tag{3.7}$$

for all vector fields X, W .

Replacing X by ξ in (3.7), we get

$$A(W) = \eta(W)\eta(\rho), \tag{3.8}$$

for any vector field W , where $A(\xi) = g(\xi, \rho) = \eta(\rho)$, ρ being the vector field associated to the 1-form A i.e., $g(X, \rho) = A(X)$.

From (3.7) and (3.8), we state the following:

Theorem 3.1. *In a projective ϕ -recurrent Kenmotsu manifold (M^{2n+1}, g) , $(n > 1)$, the characteristic vector field ξ and the vector field ρ associated to the 1-form A are co-directional and the 1-form A is given by (3.8).*

From (2.14) it follows that

$$(\nabla_w P)(X, Y)\xi = (\nabla_w R)(X, Y)\xi - \frac{1}{2n} [(\nabla_w S)(Y, \xi)X - (\nabla_w S)(X, \xi)Y]$$

Using (2.5), (2.6) and (2.9) in the above equation, we have

$$\begin{aligned} (\nabla_w P)(X, Y)\xi &= [g(W, X) - \eta(W)\eta(X)]Y - [g(W, Y) - \eta(W)\eta(Y)]X \\ &\quad - R(X, Y)(W - \eta(W)\xi) \end{aligned} \quad (3.9)$$

By virtue of (2.8), it follows from (3.9) that

$$\eta((\nabla_w P)(X, Y)\xi) = 0 \quad (3.10)$$

Using (3.9) and (3.10) in (3.1), we have

$$\begin{aligned} &-[g(W, X) - \eta(W)\eta(X)] + [g(W, Y) - \eta(W)\eta(Y)] + R(X, Y)(W - \eta(W)\xi) \quad (3.11) \\ &= A(W)[R(X, Y)\xi - (\eta(X)Y - \eta(Y)X)] \end{aligned}$$

Again using (3.8) and (2.6) in (3.11) we obtain

$$R(X, Y)W = [g(X, W)Y - g(Y, W)X] \quad (3.12)$$

Hence we can state the following:

Theorem 3.2. *A projective ϕ -recurrent Kenmotsu manifold (M^{2n+1}, g) , $(n > 1)$, is a space of constant curvature.*

We now suppose that a Kenmotsu manifold (M^{2n+1}, g) , $(n > 1)$, is projective φ -recurrent. Then from (3.3) and (3.9), it follows that

$$(\nabla_w P)(X, Y)Z = [g(Y, W)g(X, Z) - g(X, W)g(Y, Z) + g(R(X, Y)W, Z)]\xi - A(W)P(X, Y)Z \quad (3.13)$$

Next, we suppose that in a projective φ -recurrent Kenmotsu manifold, the sectional curvature of a plane $\pi \subset T_p M$ defined by

$$K_p(\pi) = g(R(X, Y)Y, X),$$

is a non-zero constant k , where $\{X, Y\}$ is any orthonormal basis of π . Then we have

$$g((\nabla_z R)(X, Y)Y, X) = 0 \quad (3.14)$$

From (2.14), we have

$$(\nabla_z P)(X, Y)Y = (\nabla_z R)(X, Y)Y - \frac{1}{2n}[(\nabla_z S)(Y, Y)X - (\nabla_z S)(X, Y)Y]$$

Using (3.14) and the above equation we have

$$g((\nabla_z P)(X, Y)Y, X) = 0 \quad (3.15)$$

By virtue of (3.15) and (3.1) we obtain

$$g((\nabla_z P)(X, Y)Y, \xi)\eta(X) = A(Z)g(P(X, Y)Y, X) \quad (3.16)$$

Since in a projective φ -recurrent Kenmotsu manifold, the relation (3.13) holds good. Using (3.13) in (3.16) and on simplification we get

$$\eta(\rho) = 0$$

Hence by (3.8) we obtain from (2.13) that

$$\varphi^2((\nabla_w P)(X, Y)Z) = 0$$

This leads to the following theorem:

Theorem 3.3. *If a projective φ -recurrent Kenmotsu manifold (M^{2n+1}, g) , $(n > 1)$, has a non-zero constant sectional curvature, then it reduces to a locally projective φ -symmetric manifold.*

References

1. Arindam Bhattacharya, "A Type of Kenmotsu Manifolds", *Ganita* 54(1)(2003), 59-62.
2. U.C. De, and G. Pathak, "On 3-Dimensional Kenmotsu Manifolds", *Indian J. Pure, Appl. Math.* 35(2) (2004), 159-165.
3. U.C. De, et al. "On φ -Recurrent Sasakian Manifold", *Novi. Sad J. Math.* 33(2) (2003), 43-48.
4. K. Kenmotsu, "A Class of Almost Contact Riemannian Manifolds", *Tohoku Math. J.*, 24 (1972), 93-103.
5. Takahashi, T., "Sasakian φ -symmetric spaces", *Tohoku Math. J.*, 29 (1977), 91-113.
6. Venkatesha and C.S. Bagewadi, "On φ -Recurrent Kenmotsu Manifolds" Communicated.
7. Venkatesha and C.S. Bagewadi, "On 3-Dimensional trans-Sasakian Manifolds", Communicated.