



BIRD DIVERSITY ON THE CAMPUS OF THE INDIAN INSTITUTE OF SCIENCE – AN EVALUATION OF TWO METHODS OF ESTIMATION

Surendra Varma*

ABSTRACT

Two methods namely, short strip transect count (SSTC) and point count (PC), were evaluated to select a robust method of estimating bird diversity on the campus of the Indian Institute of Science, a man-made ecosystem. For bird species/h (PC = 6.7, SSTC = 3.0) and the number of birds/km²h (PC = 184.7, SSTC = 40.3), the PC method encountered more species and individuals in relatively shorter time. The mean number of sightings and species per minute (sightings: PC = 1.31, SE 0.08, SSTC = 0.94, SE = 0.07, species: PC = 1.09, SE 0.06 and SSTC = 0.67, SE 0.03) were also more in the PC method and the differences are very significant (sightings: $z = 3.09$, $p < 0.001$ and for species, $z = 5.48$, $p < 0.001$). Species accumulation curve richness, diversity ($H' = 1.98$ (PC) and 1.75 (SSTC)), and evenness (E) (0.610 for PC, and 0.50 for SSTC) also favoured the PC method. These results indicate that the PC is an appropriate method for estimating bird diversity. The paper also discusses the possible ecological reasons for PC being a robust method.

Keywords: Bird diversity, species richness, bird count

* Asian Elephant Research and Conservation Centre, Centre for Ecological Sciences, Indian Institute of Science (IISc), Bangalore 560 012, India, email: varma@cesiisernetin

1. Introduction

Measures of diversity are frequently used as indicators of the well-being of ecological systems. Birds, being most diverse communities and representing a variety of habitat niches, are potentially useful as indicators of habitat changes and for other conservation-oriented approaches. Understanding such dynamic patterns of diversity is dependent on the methods employed. If there are several methods to choose from, it can sometimes be difficult to decide on the most suitable method of measuring diversity. Here, two methods of measuring bird diversity are compared, and the difference and influence of these methods on studying bird diversity evaluated in a thickly wooded, ornithologically well-known Indian Institute of Science campus. This kind of study may also help in devising strategies for preserving flora and fauna found in man-made ecosystems, which are equally important as other ecosystems, for conserving biological diversity.

2. Material and methods

2.1 Study area

The study area, the Indian Institute of Science (IISc) campus, has an area of 180 hectares and was probably an open scrubland prior to its establishment in the 1910s. The land-use pattern has undergone significant changes over the past several years. A major change has been the increase in tree cover and decrease in area under scrub and open land. The campus has a number of species of avenue trees, several species of *Cassia*, *Tabebuia* and groves of *Ficus*. Fruit-yielding trees such as *Syzygium*, *Artocarpus*, *Anona* and *Muntingia calabura* are found here. Plantations of *Acacia*, thorny *Acacia* and *Casurina*, dense thickets of *Lantana*, grasslands and open grounds also exist. A small area of grassy marsh and pool is the wetland habitat here. These microhabitats provide ideal habitats for a variety of species. More than 100 species of birds have been recorded on the campus (Pers. Obs.).

2.2 Study method

The observations of bird diversity studies were made by two methods: (i) Observers walked for 5 min continuously and recorded the bird species encountered while walking and (ii) Observers stopped for 2 min and recorded the bird species. The methods used will henceforth be referred to as short-strip transect counts (SSTC) for continuous walk method and point counts (PC) for stop method. In the SSTC method, for every five min, an average distance of 30 m was covered. All birds seen within 20 m (10 m on either side of the transect) belt were recorded. Totally,

126 five-min observations were made in the SSTC method. In the PC method, all birds seen within a 10 m radius of the stationary observer were recorded and 116 were stops made. Information such as the name of the species, the number of individuals, etc was recorded during data collection in both the methods.

The good network of roads and man-made paths in IISc was used for this study. All samplings were done between 0630 and 0830, and 1600 and 1800 hours. The data were collected during October-November 1996. There were totally eight observers, divided into four groups of two individuals each. The study area was divided into four different blocks. Every day, two blocks were selected for data collection.

2.3 Data analysis

The data were analyzed separately for both the methods. All the 126 five-min observations of the SSTC method were pooled together and species encountered per hour were calculated. In the SSTC method, the area covered was calculated by the width of the strip and the distance covered. Using this, the number of species per hour per km² was obtained. In the PC method, a 10 m radius of 116 observations was converted into km² and the number of species per hour per km² was obtained. Apart from this, all the 5 and 2 min observations of the SSTC and PC methods, respectively, were standardized into 1-min observations. Mean sightings and species per 1 min and their standard deviation (SD) were calculated. Differences were tested using the z-test. To identify the method, which gives maximum species in relatively least time, we used species and sightings per hour, per km² and per minute.

As the sample sizes of these two methods were not equal, rarefaction model for measuring species richness was used. The rarefaction method calculates the number of species expected from different communities, if all sample sizes are reduced to a standard size Program RAREFRAC BAS to compute the rarefaction curves and a sample size of $n = 500$ were used as standard. At this sample size, the SSTC and PC methods were rated in terms of their species richness. The species richness was compared based on time scale for which species accumulation curve was developed. The cumulative number of species seen after every 15 min was plotted and the relationship between time and species encountered was identified for these two methods.

Species diversity for these two methods was calculated using Shannon index. It was calculated from the equation.

$$H' = -\sum p_i \ln p_i$$

Where H' is diversity measure and the quantity p_i is the proportion of individuals found in the i th species

The t test was used to test if the values of diversity indices by the two methods are statistically significantly different. The variance in diversity of the two methods was calculated using the formula

$$\text{Var } H' = \frac{\sum p_i (\ln p_i)^2 - (\sum p_i \ln p_i)^2}{N} - \frac{S-1}{2N^2}$$

where var H' is its variance

The formula used for the t test is

$$t = \frac{H'_1 - H'_2}{(\text{Var } H'_1 + \text{Var } H'_2)^{1/2}}$$

where H'_1 is the diversity of method 1 and var H'_1 is its variance

The degree of freedom was calculated using the formula

$$df = \frac{(\text{Var } H'_1 + \text{Var } H'_2)^2}{[(\text{Var } H'_1)^2/N_1] + [(\text{Var } H'_2)^2/N_2]}$$

where N is the number of individuals

To identify how equally abundant the species measured by the SSTC and PC methods are, evenness was calculated using the formula

$$E = H' / \ln S$$

where H' is the diversity and S the number of species

3. Results

A total of 35 species (Appendix I) were observed during the study period. Table I summarizes the results of the total time spent, the area covered, the number of species, the number of sightings and the total number of birds seen for species for both SSTC and PC methods individually.

Though the time spent for the PC method was relatively lesser than that of the SSTC method (4 h compared to SSTC's 10 h), the number of species encountered

in the PC method was more than that of SSTC (6.7 species/h for PC and 3.0 species/h for SSTC). The number of individuals encountered per hour per km² was also more in the PC method (184.7 birds/km² per hour for the PC method and 40.3 birds/km² per hour for the SSTC method).

Table 1. Number of sightings and individuals for both PC and SSTC methods

SI no	Species	PC		SSTC	
		Sightings	Total number	Sightings	Total number
1	AWW	1	1	3	4
2	BD	11	12	40	64
3	BK	0	0	1	1
4	BL	0	0	2	2
5	BRP	7	16	19	53
6	BWK	1	1	3	3
7	CE	3	5	5	8
8	CM	30	273	76	981
9	GO	0	0	2	3
10	GI	2	6	4	15
11	HC	113	335	234	741
12	HO	1	1	0	0
13	HOS	0	0	1	1
14	HS	2	2	2	4
15	IR	0	0	1	1
16	KO	20	23	22	24
17	MPR	3	5	5	5
18	PFC	1	1	1	2
19	PH	4	13	15	37
20	PK	16	20	25	38
21	PRS	6	21	11	17
22	RRP	17	52	49	154
23	RVB	1	2	0	0
24	RWL	0	0	1	1
25	SD	11	17	17	22
26	SGB	5	13	13	18
27	SH	0	0	1	1
28	SMM	1	1	0	0
29	SM	0	0	1	1
30	SO	0	0	1	1
31	TB	2	4	3	3
32	IFP	31	61	64	85
33	WBK	4	5	6	6
34	WE	2	2	3	3
35	WHB	14	30	20	45

Method	Time spent (h)	No. of species	No. of individuals	Species/ (h)	Area covered (km ²)	Birds / (km ² /h)
SSTC	10.5	32	2334	3.05	0.075	40.3
PC	3.8	26	922	6.72	0.036	184.5

Standardizing the observations to mean birds per minute and comparing the results of the SSTC and PC methods it was found that the number of bird sightings and species encountered per minute was more in the PC method. The mean number of sightings per minute for the PC method was 1.31 ($n = 116$ SE 0.08) and 0.94 for the SSTC method, ($n = 126$ SE 0.07). The mean number of species per minute for the PC method was 1.09 ($n = 116$ SE 0.06) and 0.67 for the SSTC method ($n = 126$ SE 0.03). The z test shows that the differences are very significant (for sightings $z = 3.09$, $p < 001$ and for species, $z = 5.48$, $p < 001$), indicating that the PC method encountered more sightings and species of birds in a shorter period of time.

It is expected that as the time spent increases, species accumulation increases at decreasing rate and an average number of sightings and species over a large time span and area may tend to give smaller average. The smaller average for the SSTC method could be due to the same reason. For an acceptable comparison, time spend and area covered for both the methods were equalized. To equalize the time spend for SSTC, 230 min was taken and found, the mean sightings and species of birds for the SSTC method were 0.60 ($n = 46$ SE 0.06) and 0.50 ($n = 46$ SE 0.043), respectively. Apart from this, the actual averages (not standardized to 1 min observations) of both sightings and species of both the methods were compared SSTC estimated higher averages for both. However, statistically the results were not different ($z = 1.134$, $p > 0.05$ for species and $z = 0.953$, $p > 0.05$ for sightings). The area covered per hour was equalized to the PC method (0.036 km²) and the data of 62 observations of the SSTC method (area of 0.039 km²) were compared with the PC method and the SSTC method estimated 131.1 birds/km²/h. Based on these results, it can be observed that the PC method encountered more bird sightings and species and number of birds/km²/h.

Our results on species richness based on rarefaction model also favoured the PC method. Using standard sample size of 500, the expected number of species for the PC method $E(S_{500})$ was 26 and for SSTC, $E(S_{500})$ 25. The number, 26 species, could be expected only at the sample size of 600 for the SSTC method. From these results, it could be concluded that the PC method has shown the highest richness and SSTC the lowest. However, it may be noted that the difference may not be statistically significant and the species increase for the PC method was only one at a standard sample size of 500 (Fig. 1).




Fig. 1. Rarefaction curves of PC and SSTC methods showing the expected number of species as a function of sample size.

Comparing the results of species accumulation curve of both the methods, we find that the total number of species encountered in the PC method was 26, achieved in 224 min of observation, compared to 32 in a period of 570 min in SSTC method. The 26 species encountered in the PC method were observed in the SSTC method only after 330 min. of observation, needing an additional 100 min of observation in the SSTC method (Fig. 2).

The PC method also led to a higher value of species diversity. The diversity for the PC method was $H' = 1.98$, while it was 1.75 for the SSTC method. The t test revealed that the differences are highly significant (df 1745, t value 4.58, $p < .001$) in terms of diversity of birds occurring in them. Hence, the PC method has shown more diversity than the SSTC method.

The measure of evenness for both the methods showed a higher evenness for the PC method (0.610 compared to 0.504 of the SSTC method) making the PC method the most appropriate one for measuring species diversity.




Fig. 2. Species accumulation curves for both PC and SSTC methods as a function of time.

4. Discussion

By using diversity indices (Shannon index) for species diversity, it is found that the PC method gives higher diversity measure suggesting that it is an appropriate method for measuring species diversity. Although our results based on the rarefaction model for species richness show that the PC method has the highest richness, the difference may not be significant as the results are not validated by any statistical test. The species increase for the PC method was only one at the standardized sample size of 500. In the case of species evenness, it is rated between 0 and 1, value 1 representing equal abundance. In this study, the PC method estimates a higher evenness of 0.6, which is more than that of SSTC (0.5). However, the difference of species is not significantly different.

For all the three exercises, species and sightings per hour, species and sightings per hour per km² and species and sightings per minute, the PC method encountered more sightings and species of birds in a relatively shorter time. The results of species accumulation curve for both the methods also favoured the PC method. Based on this it can be concluded that more birds are sighted during the PC count and it could be a reason for the PC method to show higher species diversity.

The reasons for the PC method being an appropriate method for measuring species diversity could be that during point counts more birds could be sighted per unit time. A similar study experienced that smaller birds tend to be sighted only when the observer stands at one place and observes carefully and are probably missed while walking. For example, it was found that the number of the

sightings of white eye (*Zosterops palpebrosa*) nearly doubles in standing as compared to walking. It was also found that the birds of large body size showed no specific patterns. Some species were seen more while standing, while some were seen more while walking. Although the PC method encountered more bird sightings and species, the SSTC method has encountered more rare species than the PC method; however, the number of sightings of these rare birds (for each species) is only one. If more time is spent (it is expected that spending more time and covering larger area has the advantage of encountering more rare species), it can be assumed that those rare species also may be encountered by the PC method.

5. Conclusion

The current study was restricted to a limited period of time, as it did not cover all seasons. The seasonal changes in the diversity and their influence on the results could not be evaluated. However, as there was more manpower (with 112 man-hours), the study was designed so as to ensure an adequate sample size for measuring bird diversity. These kinds of studies are important as they provide information on the biological diversity and trend in population numbers of different species found in man-made ecosystems. Most of the well-wooded man-made ecosystems, because of rapid urbanization, are becoming increasingly isolated. This isolation has prevented the emigration of species, particularly birds from the neighboring areas. Santharam felt that the IISc campus still appears good to support a variety of species, particularly ones such as Black-rumped Flame back (*Dinopium benghalense*), Rufous Treepie (*Dendrocitta vagabunda*), Common lora (*Aegithina tiphia*) and White-browed Bulbul (*Pycnonotus luteorus*). These species existed earlier and have in recent years become locally extinct on the campus. Very interestingly, Gadagkar *et al* reported the sighting of the Black-rumped Flame back on the campus. Thus, this kind of short, quick and well-planned studies also help to assess the changes in the species diversity and their number.

Acknowledgments

Mr. I.R. Shankar Raman, Dr. N.V. Joshi, Ms. Maulishree Agrahari and Ms. Cheryl Nath of the Centre for Ecological Sciences, IISc, and Mr. P.R. Jaikrishnan examined the manuscript and suggested many useful changes. I wish to thank the members of the "Nature interest informal group" for their interest and help in collecting the data.

References

1. Magurran A.E, *Ecological diversity and its measurement*, Croom Helm Ltd, 11 New Fetter Lane, London, 1988.
2. Shyamal L, The birds of Indian Institute of Science Campus—Changes in the Avifauna *News/ Bird Watchers*, 1994, 34(1) 7-9.
3. Suresh H.S. and Bhai H.R, Flora of Indian Institute of Science Campus, Bangalore, Technical report, Centre for Ecological Sciences, Indian Institute of Science, Bangalore, India, 1998, TR No. 77.
4. Ludwig L.A. and Reynolds J.F, *Statistical ecology A primer on methods and computing*, Wiley-Interscience, 1988.
5. Varma.S, *A study of bird diversity in the Indian Institute of Science Campus, Bangalore India. A comparison of two different methods* (undated, unpublished manuscript).
6. Santharam V, Some random notes and observation on the birds of Bangalore, *News/ Bird Watchers*, 1990, 30(1), 5-7.
7. Gadagkar V *et.al*, A blue-throated flycatcher, Indian Great Reed Warbler, Common Rosewrich and Lesser Golden backed wood pecker—Four new species in the Indian Institute of Science Campus, Bangalore *News/ Bird Watchers* 1995 35(4), 69-70.

Appendix I

Birds sighted during the study period

Sl. no.	Species code	Name	Scientific name
1	AWW	Ashy Prinia	<i>Prinia socialis</i>
2	BD	Black Drongo	<i>Dicrurus macrocurcus</i>
3	BK	Brahminy Kite	<i>Haliasur Indus</i>
4	BL	Rufous-winged Bushlark	<i>Mirafra assamica</i>
5	BRP	Rock Pigeon	<i>Columba Livia</i>
6	BWK	Black-shouldered Kite	<i>Elanus caeruleus</i>
7	CE	Cattle Egret	<i>Bubulcus ibis</i>
8	CM	Common Myna	<i>Acridotheres tristis</i>
9	GO	Eurasian Golden Oriole	<i>Oriolus oriolus</i>
10	GI	Great Tit	<i>Parus major</i>
11	HC	House Crow	<i>Corvus splendens</i>
12	HO	Common Hoopoe	<i>Upupa epops</i>
13	HOS	House Sparrow	<i>Passer domesticus</i>
14	HS	House Swift	<i>Apus affinis</i>

15	IR	Indian Robin	<i>Saxicoloides fulicata</i>
16	KO	Asian Koel	<i>Eudynamys scolopacea</i>
17	MPR	Oriental Magpie-robin	<i>Copsychus saularis</i>
18	PFC	Asian Paradise Flycatcher	<i>Terpsiphone paradisi</i>
19	PH	Chinese Pond Heron	<i>Ardeola bacchus</i>
20	PK	Black Kite	<i>Milvus migrans govinda</i>
21	PRS	Purple-rumped Sunbird	<i>Nectarina zeylonica</i>
22	RRP	Rose-ringed Parakeet	<i>Psittacula cafer</i>
23	RVB	Red-vented Bulbul	<i>Vanellus Kranneri</i>
24	RWL	Red-wattled Lapwing	<i>Vanellus cafer</i>
25	SD	Spotted Dove	<i>Streptopella chinensis</i>
26	SGB	White-cheeked Barbet	<i>Megalaima virdis</i>
27	SH	Shikra	<i>Accipiter badius</i>
28	SMM	Small Minivet	<i>Pericrocotus cinnamomeus</i>
29	SM	Scaly-breasted Munia	<i>Lonchura punctulata</i>
30	SO	Spotted Owlet	<i>Athene brama</i>
31	IB	Common lailorbird	<i>Orthotomus sutorius</i>
32	IFP	Pale-billed Flowerpecker	<i>Dicaeum erythrohynchos</i>
33	WBK	White-throated Kingfisher	<i>Halcyon smyrnensis</i>
34	WE	Oriental White-eye	<i>Zosterops palpebrosa</i>
35	WHB	Yellow-billed Babbler	<i>Turdoides affinis</i>
