



On Some Structural Properties of $G_{m,n}$ Graphs

Ivy Chakrabarty*

Abstract

This is the continuation of the study on an undirected graph $G_{m,n}$ where vertex set $V = I_n = \{1, 2, 3, \dots, n\}$ and $a, b \in V$ are adjacent if and only if $a \neq b$ and $a + b$ is not divisible by m , where $m(> 1) \in \mathbb{N}$. In the present paper we computed the diameter, Wiener index, degree distance, independence number of the graph $G_{m,n}$. We also studied the complement of the graph $G_{m,n}$.

Keywords: Divisibility graph, power graph, annihilator graph, connected graph, Wiener index, degree distance, independence number of graphs

Mathematics Subject Classification (2010): 05C10

1. Wiener index and degree distance of $G_{m,n}$

Let G be a graph with vertex set $V(G)$ and edge set $E(G)$. The distance $d(u, v)$ between any two vertices $u, v \in V(G)$ is the minimum number of edges on a path in G between u and v . The diameter of a graph G is the maximum of distances between the every pairs of vertices in $V(G)$.

The Wiener index of the vertex v in G is defined as $W(v, G) = \sum_{u \in V(G)} d(u, v)$. The Wiener index of a graph G is defined as $W(G) = \sum_{\{u,v\} \subseteq V} d(u, v)$. The degree distance of a graph G is defined as $DD(G) = \sum_{\{u,v\} \subseteq V} (\deg(u) + \deg(v))d(u, v)$.

Lemma 1.1. *For any m, n , the diameter of the graph $G_{m,n}$ is less than or equal to 2.*

Proof. Let $G = G_{m,n}$. Let $a, b \in V$, where $a \neq b$ and a, b are adjacent, then $d(a, b) = 1$. Let $m \geq 2n$, then the graph $G_{m,n}$ is complete[3] so $d(i, j) = 1$ for all $i, j \in V$. Let $m < 2n$.

*Department of Mathematics, Gopalan College of Engineering and Management, Bangalore-560048, India; ivy.chakrabarty@yahoo.com

Case I. Let $m < n$, then the vertex set $V = \{1, 2, \dots, m - i, \dots, 2m - i, \dots, km - i, \dots, n\}$. Consider the set $P(x) = \{m - x, 2m - x, \dots, km - x\}$. Let $x, y \in V$, the vertex x is adjacent to the vertex y if $y \notin P(x)$. Let $y \in P(x)$, then the vertex x is not adjacent to the vertex y . Again for each $y \in V$, the vertex y is not adjacent to any other vertex $y_1 \in P(y)$ where $P(y) = \{m - y, 2m - y, \dots, km - y\}$. Consider the set $V - \{P(x) \cup P(y)\}$. Let $x, y \in V$, where $x, y \neq km$ for some $k \in \mathbb{N}$. Then the vertex $v = km \in V - \{P(x) \cup P(y)\}$. Again let $x \neq km, y = km$ for some $k \in \mathbb{N}$ then the vertex $v = y \in V - \{P(x) \cup P(y)\}$. Lastly consider $x = k_1m, y = k_2m$, where $k_1, k_2 \in \mathbb{N}$ then the vertex $v = 1 \in V - \{P(x) \cup P(y)\}$. Thus the set $V - \{P(x) \cup P(y)\}$ is nonempty. Let $z \in V - \{P(x) \cup P(y)\}$. Then the vertex z is adjacent to the vertex x as well as to the vertex y . Thus the vertices x, y are connected via the vertex z . Hence, $d(x, y) = 2$.

Case II. Let $2 < n < m < 2n$. Let $x, y \in V$. By definition, the vertex x and y are not adjacent if and only if m divides $(x + y)$. But in this case $2n < 2m$ which implies m divides $(x + y) \Leftrightarrow m = x + y$. Then the only non adjacent pairs of vertices are $P = \{(n, m - n), (n - 1, m - n + 1), \dots, (\frac{m}{2}, \frac{m}{2})\}$ if m is even and $P = \{(n, m - n), (n - 1, m - n + 1), \dots, (\frac{m+1}{2}, \frac{m-1}{2})\}$ if m is odd. Let $(x, y) \in P$, then the vertices x, y are not adjacent. Then for any $z \in V$, where $z \notin \{x, y\}$, the vertices x, y are adjacent to the vertex z . Thus the vertices x, y are connected via the vertex z , hence $d(x, y) = 2$. Again for any $(x, y) \notin P$, then the vertices x, y are adjacent.

Case III. Let $m = n$ and $a, b \in V$.

Case A. Let the vertices a and b are adjacent, then $d(a, b) = 1$.

Case B. Let the vertices a, b are not adjacent. Since $m = n$, m does not divide $(n + a)$ for all $a \in V$ which implies the vertices n and a are adjacent. Similarly we can say the vertices n and b are adjacent. So the vertices a, b are connected via the vertex n , which gives $d(a, b) = 2$. Thus we can conclude that the distance between any two distinct vertices in $G_{m,n}$ is 1 or 2. □

Theorem 1.2. Let $m \geq 2n$, then the diameter of the graph $G_{m,n}$ is one.

Proof. For $m \geq 2n$, $G_{m,n}$ is complete, hence the diameter of the graph $G_{m,n}$ is one. □

Theorem 1.3. Let $m < 2n$, then the diameter of the graph $G_{m,n}$ is two.

Proof. For $m < 2n$, the diameter of $G_{m,n}$ is two, which follows from the lemma 1.1. □

Theorem 1.4. Let $G = G_{m,n}$ be a graph. Then the Wiener index of any vertex $i \in V$, where V is the vertex set of the graph G is $W(i, G) = 2n - \text{deg } i - 2$.

Proof. Let $i \in V(G)$. Then the Wiener index of i is $W(i, G) = \sum_{j \in V(G)} d(i, j)$. The distance between any two distinct vertices in G is 1 or 2. The total number of vertices of distance 1 from i is equal to the number of

vertices adjacent to i which is nothing but the degree of the vertex i . Again the vertices which are not adjacent to the vertex i are at the distance two and the number of such vertices are $n - \deg i - 1$. Thus $W(i, G) = \sum_{j \in V(G)} d(i, j) = \deg i + (n - \deg i - 1) \cdot 2 = 2n - \deg i - 2$. \square

Theorem 1.5. *Let $m = 2$ and n be even. The the Wiener index of the graph $G_{2,n}$ is $\frac{3}{4}n^2 - n$.*

Proof. Let $m = 2$ and n be even. Let $V_1 = \{1, 3, \dots, n-1\}, V_2 = \{2, 4, \dots, n\}$ where $V = V_1 \cup V_2$. Then the degree of each vertex $i \in V_1$ and $j \in V_2$ is $\frac{n}{2}$. Let i, j be adjacent, then $d(i, j) = 1$ and $\sum_{(i,j) \subseteq V} d(i, j) = \frac{1}{2}(n \cdot \frac{n}{2} \cdot 1)$. Again consider $i, j \in V$ where i, j are not adjacent, then $d(i, j) = 2$ and each vertex is not adjacent to $n - \frac{n}{2}$ number of vertices, thus $\sum_{(i,j) \subseteq V} d(i, j) = \frac{1}{2}n \cdot (n - \frac{n}{2} - 1) \cdot 2$. Thus the Wiener index of the graph $G_{2,n}$ is

$$W(G_{2,n}) = \sum_{(i,j) \subseteq V} d(i, j) = \frac{1}{2}(n \cdot \frac{n}{2} \cdot 1) + \frac{1}{2}n \cdot (n - \frac{n}{2} - 1) \cdot 2 = \frac{3}{4}n^2 - n.$$

\square

Theorem 1.6. *Let $m = 2$ and n be even. The degree distance of $G_{m,n}$ is $\frac{1}{4}[n^2(3n-4)]$.*

Proof. Consider the graph $G = G_{2,n}$ where n be even. Let $V_1 = \{1, 3, \dots, n-1\}, V_2 = \{2, 4, \dots, n\}$ where $V = V_1 \cup V_2$. The order of the sets $V_1 = V_2 = \frac{n}{2}$. Again the degree of each vertex in V_1 and V_2 is $\frac{n}{2}$. Again no two vertices in V_1 or in V_2 are adjacent, hence the distance between any two vertices either in V_1 or in V_2 are 2. But the distance between any two vertices where $v_i \in V_1, v_j \in V_2$ is 1 as they are adjacent. Let $i \in V_1$ and $j \in V_2$ then $\sum_{(i,j) \subseteq V} (\deg(i) + \deg(j))d(i, j) = (\frac{n}{2} + \frac{n}{2}) \cdot 1 \cdot \frac{n}{2} \cdot \frac{n}{2}$.

Again let $i, j \in V_1$ or $i, j \in V_2$ then $\sum_{(i,j) \subseteq V} (\deg(i) + \deg(j))d(i, j) = (\frac{n}{2} + \frac{n}{2}) \cdot 2 \cdot (\frac{n}{2})$. Thus $\sum_{(i,j) \subseteq V} (\deg(i) + \deg(j))d(i, j) = (\frac{n}{2} + \frac{n}{2}) \cdot 1 \cdot \frac{n}{2} \cdot \frac{n}{2} + (\frac{n}{2} + \frac{n}{2}) \cdot 2 \cdot \frac{(\frac{n}{2})(\frac{n}{2}-1)}{2} + (\frac{n}{2} + \frac{n}{2}) \cdot 2 \cdot \frac{(\frac{n}{2})(\frac{n}{2}-1)}{2} = \frac{n^3}{4} + \frac{n^2(n-2)}{4} + \frac{n^2(n-2)}{4} = \frac{n^2}{4}(3n-4)$. \square

Theorem 1.7. *Let $m = 2$ and n be odd. Then the Wiener index of the graph $G_{2,n}$ is $\frac{1}{4}(n-1)(3n-1)$.*

Proof. Let $m = 2$ and n be odd. Let $V_1 = \{1, 3, \dots, n\}, V_2 = \{2, 4, \dots, n-1\}$ where $V = V_1 \cup V_2$. Then the degree of each vertex $i \in V_1$ is $\frac{n-1}{2}$ and the degree of each vertex $j \in V_2$ is $\frac{n+1}{2}$. Thus $\frac{n-1}{2}$ vertices of V_1 are at distance one from $\frac{n+1}{2}$ vertices of V_2 and vice versa. Again $(\frac{n+1}{2} - 1)$ vertices of V_2 are at distance two from $\frac{n+1}{2}$ vertices of V_2 . Similarly $(\frac{n-1}{2} - 1)$ vertices of V_1 are at distance two from $\frac{n-1}{2}$ vertices of V_1 . Thus the Wiener index of $G_{2,n}$ is $W(G_{2,n}) = \sum_{(i,j) \subseteq V} d(i, j) = \frac{1}{2}[(\frac{n-1}{2} \cdot 1 \cdot \frac{n+1}{2}) + ((\frac{n+1}{2} - 1) \cdot 2 \cdot \frac{n+1}{2}) + (\frac{n+1}{2} \cdot 1 \cdot \frac{n-1}{2}) + ((\frac{n-1}{2} - 1) \cdot 2 \cdot \frac{n-1}{2})] = \frac{1}{4}(n-1)(3n-1)$. \square

Theorem 1.8. Let $m = 2$ and n be odd. Then the degree distance of the graph $G_{2,n}$ is $\frac{n^2-1}{4}(3n-4)$.

Proof. Let $m = 2$ and n be odd. Let $V_1 = \{1, 3, \dots, n\}$, $V_2 = \{2, 4, \dots, n-1\}$ where $V = V_1 \cup V_2$. Then the degree of each vertex $i \in V_1$ is $\frac{n-1}{2}$ and the degree of each vertex $j \in V_2$ is $\frac{n+1}{2}$. Let $i \in V_1, j \in V_2$, $\sum_{(i,j) \in V} (\deg(i) + \deg(j))d(i, j) = (\frac{n-1}{2} + \frac{n+1}{2}) \cdot 1 \cdot \frac{n+1}{2} \cdot \frac{n-1}{2}$. Again let $i, j \in V_1$, then $\sum_{(i,j) \in V} (\deg(i) + \deg(j))d(i, j) = (\frac{n-1}{2} + \frac{n-1}{2}) \cdot 2 \cdot (\frac{n+1}{2})$. Similarly let $i, j \in V_2$, then $\sum_{(i,j) \in V} (d(i) + d(j))d(i, j) = (\frac{n+1}{2} + \frac{n+1}{2}) \cdot 2 \cdot (\frac{n-1}{2})$. Thus the degree distance of $G_{2,n}$ is $\sum_{(i,j) \in V} (d(i) + d(j))d(i, j) = \frac{n(n^2-1)}{4} + \frac{(n-1)(n^2-1)}{4} + \frac{(n-3)(n^2-1)}{4} = \frac{(n^2-1)(3n-4)}{4}$. \square

Theorem 1.9. Let $m(\neq 2)$ be a prime and n be multiple of m . Then the graph $G_{m,n}$ contains k vertices of degree $n-k$ and $n-k$ vertices of degree $n-k-1$ where $n = km, k \in \mathbb{N}$.

Proof. Let $m(\neq 2)$ be a prime and $n = km, k \in \mathbb{N}$. Let

$$V = \{1, 2, \dots, n\} = \{1, 2, \dots, m-1, m, \dots, 2m-1, 2m, \dots, km-1, km\}.$$

The vertices k_1m where $k_1 = 1, 2, \dots, k$ are adjacent to all other vertices except k_2m where $k_1 \neq k_2$ and $k_2 = 1, 2, \dots, k$. Thus the degree of the vertices of the form k_1m is $n - (k-1) - 1 = n - k$. Again the vertex $k_1m - i$ is not adjacent to the vertex $k_2m - j$ where $k_1 \neq k_2, k_1, k_2 = 1, 2, \dots, k, i, j = 1, 2, \dots, m$ and $i + j = m$. Thus the degree of the vertices of the form $k_1m - i$ is $n - k - 1$ (subtracting 1 because $k_1m - i$ is not adjacent to itself). \square

Theorem 1.10. Let $m(\neq 2)$ be a prime and $n = km$ where $k \in \mathbb{N}$. Then the Wiener index of the graph $G_{m,n}$ is $\frac{1}{2m}(n-1)(n+nm)$.

Proof. Let $G = G_{m,n}$, $m(\neq 2)$ be a prime and $n = km$ where $k \in \mathbb{N}$. Let $i \in V$ such that $i \neq k_1m$, where $k_1 = 1, 2, \dots, k (= \frac{n}{m})$. Then the vertex i is not adjacent to any other vertex of the form $k_1m - i$, thus $d(i, k_1m - i) = 2$. And there are $n - k = n - \frac{n}{m}$ number of vertices of the form $i \neq k_1m$. So for each vertex of the form $i \neq k_1m$ there are $k = \frac{n}{m}$ vertices which are at distance two and $(n - \frac{n}{m} - 1)$ vertices which are at distance one. Again there are $\frac{n}{m}$ number of vertices of the form $i = k_1m$ where $k_1 = 1, 2, \dots, k$. Consider $i \in V$ such that $i = k_1m$, then the vertex i is not adjacent to the vertex $j \in V$ where $i \neq j$ and $j = k_1m$. Thus in that case also $d(i, j) = 2$. And for each vertex of the form $i = k_1m$, there are $n - \frac{n}{m}$ number of vertices which are at distance one. Thus the Wiener index of $G_{m,n}$ is

$$\begin{aligned} W(G_{m,n}) &= \sum_{(i,j) \in V} d(i, j) \\ &= \frac{1}{2} \left[\left(n - \frac{n}{m} \right) \left\{ \frac{n}{m} \cdot 2 + \left(n - \frac{n}{m} - 1 \right) \cdot 1 \right\} + \frac{n}{m} \left\{ \left(\frac{n}{m} - 1 \right) \cdot 2 + \left(n - \frac{n}{m} \right) \cdot 1 \right\} \right] \quad \square \\ &= \frac{(n-1)(n+nm)}{2m}. \end{aligned}$$

Theorem 1.11. Let $m(\neq 2)$ be a prime and $n = km$ where $k \in \mathbb{N}$. The degree distance of $G_{m,n}$ is $(n - k)((n + k)(n - 2) + 1)$.

Proof. Let $G = G_{m,n}$ where $m(\neq 2)$ be a prime and $n = km$ for $k \in \mathbb{N}$. The degree distance of G is $DD(G) = \sum_{\{i,j\} \in V} (d(i) + d(j))d(i, j)$. Let $V = \{1, 2, \dots, m - 1, m, \dots, 2m - 1, 2m, \dots, km - 1, km\}$. The degree of the vertices k_1m is $n - k$ and the degree of the vertices $k_1m - i$ is $n - k - 1$. There are k vertices of degree $n - k$ and $n - k$ vertices of degree $n - k - 1$. Again for each $n - k$ vertices there are k vertices of degree $n - k$ which are at distance one, k vertices of degree $n - k - 1$ at distance two and $(n - k - k - 1)$ vertices of degree $n - k - 1$ at distance one. Similarly for each vertex k of the form k_1m , there are k vertices of degree $n - k$ at distance two and $(n - k)$ vertices of degree $n - k - 1$ at distance one. Thus the degree distance of G is

$$\begin{aligned} DD(G) &= \sum_{\{i,j\} \in V} (d(i) + d(j))d(i, j) \\ &= \frac{1}{2}(n - k)(n - k - 1 + n - k - 1)2k + (n - k - 1 + n - k).1.k \\ &\quad + (n - k - 1 + n - k - 1).1.(n - k - k - 1) \\ &\quad + k(n - k + n - k)2k + (n - k + n - k - 1)1.(n - k) \\ &= (n - k)((n + k)(n - 2) + 1). \end{aligned} \quad \square$$

Theorem 1.12. Let m, n be primes and $m = n = p$. Then $G_{m,n}$ has one vertex of degree $p - 1$ and $(p - 1)$ vertices of degree $p - 2$.

Proof. Let $m = n = p$. Let $V = \{1, 2, \dots, p\}$. Let $i(\neq p) \in V$. Then the vertex $v_i = i$ is not adjacent to the vertex $v_j = j = p - i$. Thus the degree of the vertex $v_i = i$ is $p - 2$ (as it is not adjacent to itself too). Again the vertex $m = n = p$ is adjacent to all other vertices other than itself as $p \nmid i + p$ where $i(< p) \in V$. Thus the degree of the vertex $n = p$ is $p - 1$. Hence the result follows. \square

Theorem 1.13. Let $G = G_{m,n}$ where $m = n = p$, p be a prime. Then the Wiener index of the graph G is $\frac{p^2 - 1}{2}$.

Proof. Let $m = n = p$, where p be a prime. The Wiener index of a graph G is $W(G) = \sum_{\{i,j\} \subseteq V} d(i, j)$. From the above theorem, it follows that $d(p, i) = 1$ for all $i(\neq p) \in V$. Thus $\sum_{\{p,i\} \subseteq V} d(p, i) = \deg(p)$, where $\deg(p)$ represents the degree of the vertex p . Again the vertex $i(\neq p)$ is not adjacent to two vertices one is itself and the other one is $p - i$. Thus $\sum_{\{i,j\} \subseteq V} d(i, j) = \frac{(p-2) \cdot 1 \cdot (p-1)}{2}$ where $i, j \neq p$ and i, j are adjacent. Again for each vertex $i \neq p$ there is only one vertex $p - i$ which is at distance 2, so $\sum_{\{i,j\} \subseteq V} d(i, j) = \frac{2 \cdot (p-1)}{2}$ where $i, j \neq p$ and i, j are not adjacent. Hence $W(G) = \sum_{\{i,j\} \subseteq V} d(i, j) = (p - 1) + \frac{(p-1) \cdot (p-2) \cdot 1}{2} + (p - 1) = \frac{p^2 - 1}{2}$. \square

Theorem 1.14. Let $m = n = p$, where p be a prime. Then the degree distance of $G_{m,n}$ is $(p - 1)(p^2 - p - 1)$.

Proof. Let $G = G_{m,n}$ and $m = n = p$ where p be a prime. Then the degree distance of G is $DD(G) = \sum_{\{i,j\} \subseteq V} (d(i) + d(j))d(i, j) = d(p) \cdot (d(p) + d(i)) \cdot d(p, i) + (d(i) + d(j)) \cdot 1 \cdot (p - 3) \cdot \frac{p-1}{2} + (d(i) + d(j)) \cdot 2 \cdot \frac{(p-1)}{2} = (p-1)((p-1)+(p-2)) \cdot 1 + (p-2+p-2) \cdot 1 \cdot (p-3) \cdot \frac{(p-1)}{2} + (p-2+p-2) \cdot 2 \cdot \frac{(p-1)}{2} = (p-1)(p^2-p-1)$, where p represents the vertex $n = p$ and i, j represents the vertices $i, j(\neq p) \in V$. \square

2. Complement of the graph $G_{m,n}$, independence number and independence sets of the graph $G_{m,n}$

Let the graph $\bar{G}_{m,n}$ be the complement of the graph $G_{m,n}$. Then the two distinct vertices $a, b \in \bar{G}_{m,n}$ are adjacent if m divides $(a + b)$ where the vertex set $V = \{1, 2, \dots, n\}$.

Theorem 2.1. *Let $n = m$, then the independence number of $G_{m,n}$ is 2.*

Proof. Consider the graph $\bar{G}_{m,n}$. Let $V = \{1, 2, \dots, n\}$. The pair of vertices $(i, n-i)$ where $i \in V, i \neq n$ and $i \neq \frac{n}{2}$ forms cliques in $\bar{G}_{m,n}$. Thus the independent sets of $G_{m,n}$ are $\{i, n-i\}$. Hence the independence number of $G_{m,n}$ is 2 which is the cardinality of the set $\{i, n-i\}$. \square

Theorem 2.2. *Let $m = n$ where m is odd. Then the number of independent sets of $G_{m,n}$ is $\lfloor \frac{n}{2} \rfloor$.*

Proof. Consider the graph $\bar{G} = \bar{G}_{m,n}$, where $m = n$ and m is odd. Let the vertex set $V = \{1, 2, \dots, n\}$. Then the vertex $v = n$ is isolated vertex in the graph \bar{G} as $m \nmid i + n$ for $i(i \neq n) \in V$ since $i < n$. And for any vertex $i \in V$ where $i = 1, 2, \dots, n-1$ is adjacent to the vertex $n-i$ and the number of such pairs is $\lfloor \frac{n}{2} \rfloor$. Thus the number of cliques in \bar{G} is $\lfloor \frac{n}{2} \rfloor$. Hence the number of independence sets of $G_{m,n}$ is $\lfloor \frac{n}{2} \rfloor$. \square

Theorem 2.3. *Let $m = n$ where m is even. Then the number of independent sets of $G_{m,n}$ is $\frac{n}{2} - 1$.*

Proof. Consider the graph $\bar{G}_{m,n}$ where $m = n$ and m be even. Let the vertex set $V = \{1, 2, \dots, n\}$. The vertices n and $\frac{n}{2}$ are not adjacent as $m \nmid n + \frac{n}{2}$. Thus the vertices n and $\frac{n}{2}$ will not form a clique in the graph $\bar{G}_{m,n}$. Again let $j \in V$ where $j \neq n, \frac{n}{2}$. Consider the vertex $i = n-j$ where $j = 1, 2, \dots, \frac{n}{2} - 1$. Then the vertices $j, n-j$ forms cliques in $\bar{G}_{m,n}$ for all j as m divides $j + (n-j)$. Thus the number of cliques in $\bar{G}_{m,n}$ is $\frac{n}{2} - 1$. Hence the number of independent sets of $G_{m,n}$ is $\frac{n}{2} - 1$. \square

Theorem 2.4. *Let $m > n$. Then the independence number of $G_{m,n}$ is 2.*

Proof. Consider the graph $\bar{G}_{m,n}$ where $m > n$ and the vertex set $V = \{1, 2, \dots, n\}$. Then the vertices n and $m - n$ form a clique in $\bar{G}_{m,n}$. Thus the independence number of $G_{m,n}$ is 2 for $m > n$. \square

Theorem 2.5. *Let $m < n$ where $m \neq 2$. Then the independence number of $G_{m,n}$ is $\lfloor \frac{n}{m} \rfloor$.*

Proof. Let $m < n$ where $m \neq 2$. The vertices $\{m, 2m, \dots, km\}$ where $km \leq n$ forms an independent set in $G_{m,n}$ as they form a clique in $\bar{G}_{m,n}$. And the cardinality of the set $\{m, 2m, \dots, km\}$ is $\lfloor \frac{n}{m} \rfloor$. Hence the results follows. \square

Theorem 2.6. *Let $m = 2$ and $n \in \mathbb{N}$. Then the independence number of $G_{m,n}$ is $\lfloor \frac{n}{2} \rfloor$. The number of independent set is 2.*

Proof. Let $G = G_{m,n}$ where $m = 2$ and $n \in \mathbb{N}$. Let $V = \{1, 2, \dots, n\}$. The set $E_1 = \{2, 4, \dots\} \subseteq V$ form the independent set of G as no two vertices of E_1 are adjacent in G . This set is maximal. Since for a given n there are $\lfloor \frac{n}{2} \rfloor$ number of even numbers in the set $\{1, 2, \dots, n\}$. Thus the independence number of G is $\lfloor \frac{n}{2} \rfloor$. Thus the sets $O_1 = \{1, 3, 5, \dots\}$ and $E_1 = \{2, 4, 6, \dots\}$ are the independent sets of G as no two vertices in E_1 are adjacent as well as no two vertices in O_1 are adjacent. Thus the number of independent set is two. \square

3. Conclusion

In this article, we computed the diameter, Wiener index of a vertex, and Wiener index and degree distance of the graphs $G_{2,n}$, $G_{m,n}$, where $m \neq 2$ is a prime, n is a multiple of m and $G_{p,p}$, where p is a prime. In future one can study various energies, domination, planarity etc. of the graph $G_{m,n}$.

Acknowledgments

The author would like to thank Department of Mathematics, CHRIST (Deemed to be University) for providing the support and exposure in various workshops and conferences. The author is deeply grateful to Prof. Joseph Varghese for his guidance and support.

References

- [1] I. Chakrabarty, S.Ghosh and M.K.Sen, *Undirected power graphs of semigroups*, Semigroup Forum **78** (2009), 410-426.

- [2] I. Chakrabarty, S.Ghosh, T.K.Mukherjee and M.K.Sen, *Intersection graphs of ideals of rings*, Discrete Mathematics **309** (2009), 5381-5392.
- [3] I. Chakrabarty, *An undirected graph on finite subset of natural numbers*, Indian Journal of Discrete Mathematics **2 (1)** (2015), 128-138.
- [4] A. A. Dobrynin, A. A. Kochetova, *Degree distance of a graph: a degree analogue of the Wiener index*, J.Chem.Inf.Comput.Sci. **34**(1994), 1082-1086.
- [5] D. Gray, H. Wang, *Cycles, the degree distance and the Wiener index*, Open Journal of Discrete Mathematics **2** (2012), 156-159.
- [6] I. Gutman, *Selected properties of the Schultz molecular topological index*, J.Chem.Inf.Comput.Sci. **34**(1994), 1087-1089.
- [7] A. V. Kelarev and S. J. Quinn, *Directed graph and combinatorial properties of semigroups*, J. Algebra **251** (2002), 16-26.
- [8] D. B. West, *Introduction to Graph Theory*, 2nd edition, Prentice Hall, Upper Saddle River, NJ, 2001.
- [9] H. Wiener, *Structural Determination of Paraffin Boiling Points*, Journal of the American Chemical Society **69** (1947), no. 1, 17-20.
- [10] B. Zelinka, *Intersection graphs of finite abelian groups*, Czech. Math. J. **25 (100)** (1975), no. 2, 171-174.