



## Sum Labeling for Some Star and Cycle Related Special Graphs

J Gerard Rozario,\* J Jon Arockiaraj† and P Lawrence Rozario Raj‡

### Abstract

A *sum labeling* is a mapping  $\lambda$  from the vertices of  $G$  into the positive integers such that, for any two vertices  $u, v \in V(G)$  with labels  $\lambda(u)$  and  $\lambda(v)$ , respectively,  $(uv)$  is an edge iff  $\lambda(u) + \lambda(v)$  is the label of another vertex in  $V(G)$ . Any graph supporting such a labeling is called a *sum graph*. It is necessary to add (as a disjoint union) a component to sum label a graph. This disconnected component is a set of isolated vertices known as *isolates* and the labeling scheme that requires the fewest isolates is termed *optimal*. The number of isolates required for a graph to support a sum labeling is known as the *sum number* of the graph. In this paper, we will obtain *optimal sum labeling scheme* for some star and cycle related special graphs.

**Keywords:** Sum labeling, sum number, sum graph, path union, arbitrary supersubdivision

**AMS Subject Classification (2010):** 05C78

---

\* PG and Research Department of Mathematics, St Joseph's College of Arts and Science, Cuddalore, Tamil Nadu, India  
rozario.gerard@yahoo.com.

† PG and Research Department of Mathematics, St Joseph's College of Arts and Science, Cuddalore, Tamil Nadu, India  
jonarockiaraj@gmail.com

‡ Department of Mathematics, St Joseph's College, Trichirappalli, Tamil Nadu, India, lawraj2006@yahoo.co.in

## 1. Introduction

All the graphs considered here are simple, finite and undirected. For all terminologies and notations we follow Harary [1] and graph labeling as in [2]. Sum labeling of graphs was introduced by Harary [3] in 1990. Following definitions are useful for the present study.

**Definition 1.1** A *Sum Labeling* is a mapping  $\lambda$  from the vertices of  $G$  into the positive integers such that, for any two vertices  $u, v \in V(G)$  with labels  $\lambda(u)$  and  $\lambda(v)$ , respectively,  $(uv)$  is an edge iff  $\lambda(u) + \lambda(v)$  is the label of another vertex in  $V(G)$ . Any graph supporting such a labeling is called a *Sum Graph*.

**Definition 1.2** It is necessary to add (as a disjoint union) a component to sum label a graph. This disconnected component is a set of isolated vertices known as *Isolates* and the labeling scheme that requires the fewest isolates is termed *Optimal*.

**Definition 1.3** The number of isolates required for a graph  $G$  to support a sum labeling is known as the *Sum Number* of the graph. It is denoted as  $\sigma(G)$ .

**Definition 1.4** Let  $G$  be a graph with  $q$  edges. A graph  $H$  is called a *Super subdivision* of  $G$  if  $H$  is obtained from  $G$  by replacing every edge  $e_i$  of  $G$  by a complete bipartite graph  $K_{2,m_i}$  for some  $m_i, 1 \leq i \leq q$  in such a way that the end vertices of each  $e_i$  are identified with the two vertices of 2-vertices part of  $K_{2,m_i}$  after removing the edge  $e_i$  from graph  $G$ . If  $m_i$  is varying arbitrarily for each edge  $e_i$  then super subdivision is called arbitrary super subdivision of  $G$ .

**Definition 1.5** A *caterpillar* is a graph which has the property that if we remove all the vertices of degree 1 then what remains is a path. Such a path is called the *spine* of the caterpillar. The two end points of a spine are the *head* and *tail* respectively. Other vertices of the spine are called the *internal vertices*. The vertices of degree 1 of a caterpillar, other than head and tail, will be called the *feet*. These vertices are attached to the internal vertices of spine with edge called the *legs* of the caterpillar.

**Definition 1.6** A *shrub* is a tree which has at most one inner vertex. This special vertex is called the root of the shrub. All neighbours of this vertex are leaves or near leaves.

**Definition 1.7** (Chung et. al [4]) A tree is called a *spider* if it has a center vertex  $c$  of degree  $k > 1$  and each other vertex either is a leaf (pendent vertex) or has degree 2. Thus a spider is an amalgamation of  $k$  paths with various lengths. If it has  $x_1$  paths of length  $a_1$ ,  $x_2$  paths of length  $a_2$ , ...,  $x_n$  paths of length  $a_m$ , we denote the spider by  $SP(a_1^{x_1}, a_2^{x_2}, \dots, a_m^{x_m})$  where  $a_1 < a_2 < \dots < a_m$  and  $x_1 + x_2 + \dots + x_n = k$

**Definition 1.8** (Shee and Ho. et al [5]) Let  $G_1, G_2, \dots, G_n, n \geq 2$  be  $n$  copies of a fixed graph  $G$ . The graph obtained by adding an edge between  $G_i$  and  $G_{i+1}$  for  $i=1, 2, \dots, n-1$  is called *path union* of  $G$ .

## 2. Optimal Sum Labeling Scheme for Arbitrary Super Subdivision of Star Related Graphs

Sethuraman et. al. [6], introduced a new method of construction called Supersubdivision of graph and proved that arbitrary supersubdivision of any path and cycle  $C_n$  are graceful. Kathiresan et.al [7], proved that arbitrary supersubdivision of any star is graceful. In [8], Gerard Rozario et.al proved that arbitrary super subdivision of path, cycle and star are sum graph with sum number 2.

In this section, we prove that graphs obtained by arbitrary super subdivision of comb, caterpillar, shrubs and path union of spider are sum graph with sum number 2.

**Theorem 2.1** Arbitrary supersubdivision of comb  $P_n \odot K_1$  is sum graph with sum number 2.

**Proof:** Let  $G$  be a comb  $P_n \odot K_1$ . Let  $v_i (1 \leq i \leq n)$  be the vertices of path and  $w_i$  be the pendent vertex adjacent to  $v_i (1 \leq i \leq n)$ . Let  $H$  be the arbitrary supersubdivision of  $G$  which is obtained by replacing every edge of  $G$  with  $K_{2, m_i}$ . Let  $m = \sum_{i=1}^{n-1} m_i$ . Let  $u_j$  be the vertices which are used for arbitrary supersubdivision of  $G$  where  $1 \leq j \leq m$ . Let  $x$  and  $y$  be two isolated vertices. Therefore, the vertex set of  $H$  is  $V(H) = \{ v_1, v_2, \dots, v_n, w_1, w_2, \dots, w_n, u_1, u_2, \dots, u_m \}$ .

Define  $f : V(H) \rightarrow N$

$$f(v_1) = 1 ; \quad f(v_i) = f(v_{(i-1)}) + 2 \quad ; 2 \leq i \leq n$$

$$f(w_1) = 2 ; f(w_i) = f(w_{(i-1)}) + 2 ; 2 \leq i \leq n$$

$$f(u_1) = m + n \quad f(u_j) = f(u_{j-1}) - 1 ; 2 \leq j \leq m$$

Then  $f(x) = f(u_1) + 1$  and  $f(y) = f(u_1) + 2$

Thus, arbitrary supersubdivision of comb  $P_n \odot K_1$  is sum graph with sum number 2.

**Theorem 2.2** Arbitrary supersubdivision of caterpillar is sum graph with sum number 2.

Proof: Let G be a caterpillar with n vertices. Let  $v_i$  ( $1 \leq i \leq n$ ) be the vertices of G. First we need to identify the vertices of G in order to super subdivide G. We present an algorithm to identify the vertices of caterpillar. The algorithm starts from the head of the caterpillar (1<sup>st</sup> vertex of the spine). Whenever the algorithm finds a new internal vertex of the spine having degree  $\geq 3$ , it first visits all vertices of degree 1 adjacent to it before proceeding to the next vertex of spine. The head of the caterpillar is named as  $v_1$  and every new vertex identified by the algorithm is named as  $v_2, v_3, \dots, v_n$  respectively.

Let H be the arbitrary supersubdivision of caterpillar G which is obtained by replacing every edge of G with  $K_{2,m_i}$ . Let  $m = \sum_1^{n-1} m_i$ . Let  $u_j$  be the vertices which are used for arbitrary supersubdivision of G where  $1 \leq j \leq m$ . Let x and y be two isolated vertices. Therefore, the vertex set of H is  $V(H) = \{ v_1, v_2, \dots, v_n, u_1, u_2, \dots, u_m \}$ .

Define  $f : V(H) \rightarrow N$

$$f(v_i) = i ; 1 \leq i \leq n$$

$$f(u_1) = m + n \quad f(u_j) = f(u_{j-1}) - 1 ; 2 \leq j \leq m$$

Then  $f(x) = f(u_1) + 1$  and  $f(y) = f(u_1) + 2$

Hence, arbitrary supersubdivision of caterpillar is sum graph with sum number 2.

Illustration: Sum labeling for arbitrary supersubdivision of caterpillar is given in figure 2.1




Figure 2.1

**Theorem 2.3** Arbitrary supersubdivision of shrub is optimal summable with sum number 2.

*Proof:* Let  $G$  be a shrub with root  $r$  and  $b_1, b_2, \dots, b_k$  be the vertices adjacent to  $r$ . Let  $d_i = \text{deg}(b_i) - 1; i = 1, 2, \dots, k$ . Let  $c_{ij}$  be the vertices adjacent to  $b_i$  other than  $r$  for  $i = 1, 2, \dots, k$  and  $j = 1, 2, \dots, d_i$ . In order to identify the vertices of shrub we follow the following algorithm. The algorithm starts from the root of the shrub and then it identifies an adjacent vertex (say  $b_1$ ) of  $r$ . Before it proceeds to the next adjacent vertex of  $r$ , the algorithm first visits all neighbour vertices  $c_{ij}$  of  $b_1$  with degree 1 (if any) and proceeds to branches of  $c_{ij}$  of higher order (if any). The algorithm renames the root  $r$  as  $v_1$  and assigns  $v_2, v_3, \dots, v_n$  respectively to the new vertex it visits first. Now, the vertices of shrubs are renamed as  $v_1, v_2, v_3, \dots, v_n$  where  $n$  is the total number of vertices of the shrub.

Let  $H$  be the arbitrary supersubdivision of shrub  $G$  which is obtained by replacing every edge of  $G$  with  $K_{2,m_i}$ . Let  $m = \sum_1^{n-1} m_i$ . Let  $u_j$  be the vertices which are used for arbitrary supersubdivision of  $G$  where  $1 \leq j \leq m$ . Let  $x$  and  $y$  be two isolated vertices. Therefore, the vertex set of  $H$  is  $V(H) = \{v_1, v_2, \dots, v_n, u_1, u_2, \dots, u_m\}$ .

Define  $f : V(H) \rightarrow \mathbb{N}$

$$f(v_i) = i \quad ; 1 \leq i \leq n;$$

$$f(u_1) = m + n \quad ; \quad f(u_j) = f(u_{j-1}) - 1 \quad ; 2 \leq j \leq m.$$

Then  $f(x) = f(u_1) + 1$  and  $f(y) = f(u_1) + 2$

Hence, arbitrary supersubdivision of shrub is optimal summable with sum number 2.

Illustration: Sum labeling for arbitrary supersubdivision of shrub is given in figure 2.2




Figure 2.2

**Theorem 2.4** Arbitrary supersubdivision of path union of spider is sum graph with sum number 2.

**Proof:** Let  $G$  be the path union of spiders obtained from  $n$  spiders (not necessarily unique) by adding an edge between centre vertex of  $i^{th}$  spider and  $(i+1)^{th}$  spider for  $i = 1, 2, \dots, (n-1)$ . Let  $c_1, c_2, \dots, c_n$  be the center vertices of the spider. Let  $k_1, k_2, \dots, k_n$  be the number of vertices in  $i^{th}$  spider excluding the center vertex where  $1 \leq i \leq n$ . Let  $p_{11}, p_{12}, \dots, p_{1k_1}, p_{21}, p_{22}, \dots, p_{2k_2}, \dots, p_{n1}, p_{n2}, \dots, p_{nk_n}$  be the vertices of the  $i^{th}$  spider respectively. Therefore,  $V(G) = \{ c_1, c_2, \dots, c_n, p_{11}, p_{12}, \dots, p_{1k_1}, p_{21}, p_{22}, \dots, p_{2k_2}, \dots, p_{n1}, p_{n2}, \dots, p_{nk_n} \}$ . Let  $t$  be the total number of vertices of  $G$ . i.e.,  $t = n + k_1 + k_2 + \dots + k_n$ . Let  $G^*$  be the arbitrary supersubdivision of  $G$  which is obtained by

replacing every edge of  $G$  with  $K_{2,m_i}$ . Let  $m = \sum_1^{n-1} m_i$ . Let  $u_j$  be the vertices which are used for arbitrary supersubdivision of  $G$  where  $1 \leq j \leq m$ . Let  $x$  and  $y$  be two isolated vertices. Therefore, the vertex set of  $G^*$  is

$$V(G^*) = \{c_1, c_2, \dots, c_n, p_{11}, p_{12}, \dots, p_{1k_1}, \dots, p_{n1}, p_{n2}, \dots, p_{nk_n}, u_1, u_2, \dots, u_m\}.$$

Define  $f : V(G^*) \rightarrow \mathbb{N}$

$$f(c_1) = 1 \quad ; f(c_i) = f(c_{i-1}) + k_{(i-1)} + 1 \quad \text{for } 2 \leq i \leq n$$

for  $1 \leq i \leq n$

$$\begin{cases} f(p_{i1}) = f(p_1) + 1 \\ \text{for } 2 \leq j \leq k_i \\ \{ f(p_{ij}) = f(p_{i(j-1)}) + 1 \end{cases}$$

$$f(u_1) = m + t \quad f(u_j) = f(u_{j-1}) - 1 \quad ; 2 \leq j \leq m$$

Then  $f(x) = f(u_1) + 1$  and  $f(y) = f(u_1) + 2$

Hence, arbitrary supersubdivision of path union of spiders is optimal summable with sum number 2.

Illustration: Sum labeling for arbitrary supersubdivision of path union of spiders is given in figure 2.3




Figure 2.3

### 3. Optimal Sum Labeling Scheme for Cycle Related Graphs

In [9], Ponraj et.al proved that the graph  $G$  with  $V(G) = V(C_n) \cup \{v\}$  and  $E(G) = E(C_n) \cup \{c_1v, c_3v\}$  and the graph  $[C_n, P_m]$  are pair sum graph. In this section we prove that the above said two graphs are sum graphs.

**Theorem 3.1** Let  $G$  be a graph with  $V(G) = V(C_n) \cup \{v\}$  and  $E(G) = E(C_n) \cup \{c_1v, c_3v\}$ . Then  $G$  is a sum graph with sum number 2.

Proof: Let  $c_1, c_2, c_3, \dots, c_n$  be the vertices of the cycle. Let  $x$  and  $y$  be two isolated vertices. The vertex set of  $G$  is given by  $V(G) = \{c_1, c_2, c_3, \dots, c_n, v, x, y\}$

Define  $f : V(G) \rightarrow \mathbb{N}$

**Case (i) :  $n$  is odd number**

**Case (i) (a):  $n = 3$**

$$\text{For } C_3, f(c_1) = 1; f(c_2) = 2; f(c_3) = 3$$

$$f(v) = f(c_1) + f(c_3)$$

$$f(x) = f(c_1) + f(v) \text{ and } f(y) = f(c_3) + f(v)$$

Thus,  $G$  is sum graph with sum number 2 when  $n = 3$ .

**Case (i) (b):  $n = 5, 7$**

$$\text{For } C_5, C_7 \quad f(c_1) = 2; f(c_2) = 1; f(c_3) = 3$$

$$f(c_n) = f(c_1) + f(c_3)$$

$$f(c_i) = f(c_{i-1}) + f(c_{i-2}) \quad ; 4 \leq i \leq (n - 1)$$

$$f(v) = \begin{cases} f(c_n) + f(c_{n-1}) & \text{if } f(c_n) < f(c_{n-1}) \\ f(c_{n-1}) + f(c_{n-2}) & \text{if } f(c_n) > f(c_{n-1}) \end{cases}$$

$$f(x) = f(c_1) + f(v) \text{ and } f(y) = f(c_3) + f(v)$$

Thus,  $G$  is sum graph with sum number 2 when  $n = 5, 7$ .

**Case (i) (c):  $n \geq 9$**

$$\text{For } C_n, \quad f\left(c\left(\frac{n-3}{2}\right)\right) = 1; \quad f\left(c\left(\frac{n-3}{2}-1\right)\right) = 2; \quad f\left(c\left(\frac{n-3}{2}+1\right)\right) = 3;$$



$$f\left(c\left(\frac{n-3}{2}-2\right)\right) = 5; f\left(c\left(\frac{n-3}{2}+2\right)\right) = 4$$

for  $1 \leq i \leq \frac{n-9}{2}$

$$\left\{ \begin{array}{l} f\left(c\left(\frac{n-3}{2}+i+2\right)\right) = f\left(c\left(\frac{n-3}{2}-i\right)\right) + f\left(c\left(\frac{n-3}{2}-i-1\right)\right) \\ f\left(c\left(\frac{n-3}{2}-i-2\right)\right) = f\left(c\left(\frac{n-3}{2}+i+1\right)\right) + f\left(c\left(\frac{n-3}{2}+i+2\right)\right) \end{array} \right\}$$

$$f(c_{n-3}) = f(c_1) + f(c_2); f(c_n) = f(c_{n-3}) + f(c_{n-4})$$

$$f(c_{n-2}) = f(c_1) + f(c_n); f(c_{n-1}) = f(c_{n-3}) + f(c_{n-2})$$

$$f(v) = f(c_n) + f(c_{n-1})$$

$$f(x) = f(c_1) + f(v) \text{ and } f(y) = f(c_3) + f(v)$$

Thus, G is sum graph with sum number 2 when  $n \geq 9$

**Case (ii): n is even number**

**Case (ii) (a): n = 4**

For  $C_4$ ,  $f(c_1) = 1; f(c_2) = 2; f(c_4) = 3; f(c_3) = f(c_1) + f(c_4)$

$$f(v) = f(c_2) + f(c_3)$$

$$f(x) = f(c_1) + f(v) \text{ and } f(y) = f(c_3) + f(v)$$

Thus, G is sum graph with sum number 2 when  $n = 4$ .

**Case (ii) (b): n ≥ 6**

For  $C_n$ ,  $f(c_1) = 1; f(c_2) = 2; f(c_n) = 3; f(c_3) = f(c_2) + f(c_n)$

$$f(c_{n-1}) = f(c_1) + f(c_n)$$

for  $4 \leq i \leq n/2$

$$\left\{ \begin{array}{l} \text{If } i \text{ is even, } \left\{ \begin{array}{l} f(c_i) = f(c_{i-1}) + f(c_{i-2}) - 1 \\ f(c_{i+(n-2i+2)}) = f(c_{i+(n-2i+2)+1}) + f(c_{i+(n-2i+2)+2}) \end{array} \right\} \\ \text{If } i \text{ is odd, } \left\{ \begin{array}{l} f(c_i) = f(c_{i-1}) + f(c_{i-2}) \\ f(c_{i+(n-2i+2)}) = f(c_{i+(n-2i+2)+1}) + f(c_{i+(n-2i+2)+2}) - 1 \end{array} \right\} \end{array} \right\}$$

$$f\left(c_{\left(\frac{n}{2}+1\right)}\right) = f\left(c_{\left(\frac{n}{2}\right)}\right) + f\left(c_{\left(\frac{n}{2}-1\right)}\right)$$

$$f(v) = \left\{ \begin{array}{l} f\left(c_{\left(\frac{n}{2}\right)}\right) + f\left(c_{\left(\frac{n}{2}+1\right)}\right) \text{ if } f\left(c_{\left(\frac{n}{2}\right)}\right) < f\left(c_{\left(\frac{n}{2}+2\right)}\right) \\ f\left(c_{\left(\frac{n}{2}+1\right)}\right) + f\left(c_{\left(\frac{n}{2}+2\right)}\right) \text{ if } f\left(c_{\left(\frac{n}{2}\right)}\right) > f\left(c_{\left(\frac{n}{2}+2\right)}\right) \end{array} \right\}$$

$$f(x) = f(c_1) + f(v) \text{ and } f(y) = f(c_3) + f(v)$$

Thus, G is sum graph with sum number 2 when  $n \geq 6$

Hence, G is sum graph with sum number 2 for all  $n \geq 3$ .

Illustration: Sum labeling of G with  $n = 5$  and  $n = 8$  is given in figure 3.1




Figure 3.1

*Notation:* Two copies of the cycle  $C_n$  connected by the path  $P_m$  is denoted by  $[C_n, P_m]$

**Theorem 3.2** The graph  $[C_n, P_m]$  is a sum graph with sum number 2.

*Proof:* Let  $c_1, c_2, c_3, \dots, c_n$  be the vertices of the first copy of cycle  $C_n$  and  $v_1, v_2, v_3, \dots, v_n$  be the vertices of the second copy of cycle  $C_p$ . Let  $p_1, p_2, \dots, p_m$  be the vertices of path connecting the two copies of the cycle. Let  $x$  and  $y$  be two isolated vertices.

Define  $f : V(G) \rightarrow \mathbb{N}$

**Case (i):  $n$  is odd number**

**Case (i) (a):  $n = 3$**

For  $C_3$ ,  $f(c_1) = 1; f(c_2) = 2; f(c_3) = 3$

$$f(p_1) = f(c_1) + f(c_3) \quad ; \quad f(p_2) = f(p_1) + f(c_1)$$

$$f(p_i) = f(p_{(i-1)}) + f(p_{(i-2)}) \text{ for } 3 \leq i \leq m$$

$$f(v_1) = f(p_m) + f(p_{(m-1)}) \quad f(v_2) = f(v_1) +$$

$f(p_m)$

$$f(v_i) = f(v_{(i-1)}) + f(v_{(i-2)}) \text{ for } 3 \leq i \leq n$$

$$f(x) = f(v_n) + f(v_1) \text{ and } f(y) = f(v_n) + f(v_{(n-1)})$$

Thus,  $G$  is sum graph with sum number 2 when  $n = 3$ .

**Case (i) (b):  $n = 5, 7$**

For  $C_5, C_7$      $f(c_1) = 2; f(c_2) = 1; f(c_3) = 3$

$$f(c_n) = f(c_1) + f(c_3)$$

$$f(c_i) = f(c_{(i-1)}) + f(c_{(i-2)}) \quad ; 4 \leq i \leq n - 1$$

$$f(p_1) = \begin{cases} f(c_n) + f(c_{(n-1)}) & \text{if } f(c_n) < f(c_{(n-1)}) \\ f(c_{(n-1)}) + f(c_{(n-2)}) & \text{if } f(c_n) > f(c_{(n-1)}) \end{cases}$$

$$f(p_2) = f(p_1) + f(c_1)$$

$$f(p_i) = f(p_{(i-1)}) + f(p_{(i-2)}) \text{ for } 3 \leq i \leq m$$

$$f(v_1) = f(p_m) + f(p_{(m-1)})$$

$$f(v_2) = f(v_1) + f(p_m)$$

$$f(v_i) = f(v_{(i-1)}) + f(v_{(i-2)}) \text{ for } 3 \leq i \leq n$$

$$f(x) = f(v_n) + f(v_1) \text{ and } f(y) = f(v_n) + f(v_{(n-1)})$$

Thus,  $G$  is sum graph with sum number 2 when  $n = 5, 7$ .

**Case (i) (c):  $n \geq 9$**

For  $C_n$ ,  $f\left(c\left(\frac{n-3}{2}\right)\right) = 1$ ;  $f\left(c\left(\frac{n-3}{2}-1\right)\right) = 2$ ;  $f\left(c\left(\frac{n-3}{2}+1\right)\right) = 3$ ;  
 $f\left(c\left(\frac{n-3}{2}-2\right)\right) = 5$ ;  $f\left(c\left(\frac{n-3}{2}+2\right)\right) = 4$   
 for  $1 \leq i \leq \frac{n-9}{2}$

$$\left\{ \begin{array}{l} f\left(c\left(\frac{n-3}{2}+i+2\right)\right) = f\left(c\left(\frac{n-3}{2}-i\right)\right) + f\left(c\left(\frac{n-3}{2}-i-1\right)\right) \\ f\left(c\left(\frac{n-3}{2}-i-2\right)\right) = f\left(c\left(\frac{n-3}{2}+i+1\right)\right) + f\left(c\left(\frac{n-3}{2}+i+2\right)\right) \end{array} \right\}$$

$f(c_{n-3}) = f(c_1) + f(c_2)$ ;  $f(c_n) = f(c_{n-3}) + f(c_{n-4})$   
 $f(c_{n-2}) = f(c_1) + f(c_n)$ ;  $f(c_{n-1}) = f(c_{n-3}) + f(c_{n-2})$   
 $f(p_1) = f(c_n) + f(c_{n-1})$   
 $f(p_2) = f(p_1) + f(c_1)$   
 $f(p_i) = f(p_{(i-1)}) + f(p_{(i-2)})$  for  $3 \leq i \leq m$   
 $f(v_1) = f(p_m) + f(p_{(m-1)})$   $f(v_2) = f(v_1) + f(p_m)$   
 $f(v_i) = f(v_{(i-1)}) + f(v_{(i-2)})$  for  $3 \leq i \leq n$   
 $f(x) = f(v_n) + f(v_1)$  and  $f(y) = f(v_n) + f(v_{(n-1)})$

Thus, G is sum graph with sum number 2 when  $n \geq 9$

**Case (ii):  $n$  is even number**

**Case (ii) (a):  $n = 4$**

For  $C_4$ ,  $f(c_1) = 1$ ;  $f(c_2) = 2$ ;  $f(c_4) = 3$ ;  $f(c_3) = f(c_1) + f(c_4)$   
 $f(p_1) = f(c_2) + f(c_3)$ ;  $f(p_2) = f(p_1) + f(c_1)$   
 $f(p_i) = f(p_{(i-1)}) + f(p_{(i-2)})$  for  $3 \leq i \leq m$   
 $f(v_1) = f(p_m) + f(p_{(m-1)})$ ;  $f(v_2) = f(v_1) + f(p_m)$   
 $f(v_i) = f(v_{(i-1)}) + f(v_{(i-2)})$  for  $3 \leq i \leq n$   
 $f(x) = f(v_n) + f(v_1)$  and  $f(y) = f(v_n) + f(v_{(n-1)})$

Thus, G is sum graph with sum number 2 when  $n = 4$ .

**Case (ii) (b):  $n \geq 6$**

For  $C_n$ ,

$$f(c_1) = 1; f(c_2) = 2; f(c_n) = 3; f(c_3) = f(c_2) + f(c_n)$$

$$f(c_{n-1}) = f(c_1) + f(c_n) \text{ for } 4 \leq i \leq n/2$$

$$\left\{ \begin{array}{l} \text{If } i \text{ is even, } \left\{ \begin{array}{l} f(c_i) = f(c_{i-1}) + f(c_{i-2}) - 1 \\ f(c_{i+(n-2i+2)}) = f(c_{i+(n-2i+2)+1}) + f(c_{i+(n-2i+2)+2}) \end{array} \right\} \\ \text{If } i \text{ is odd, } \left\{ \begin{array}{l} f(c_i) = f(c_{i-1}) + f(c_{i-2}) \\ f(c_{i+(n-2i+2)}) = f(c_{i+(n-2i+2)+1}) + f(c_{i+(n-2i+2)+2}) - 1 \end{array} \right\} \end{array} \right\}$$

$$f\left(c_{\left(\frac{n}{2}+1\right)}\right) = f\left(c_{\left(\frac{n}{2}\right)}\right) + f\left(c_{\left(\frac{n}{2}-1\right)}\right)$$

$$f(p_1) = \left\{ \begin{array}{l} f\left(c_{\left(\frac{n}{2}\right)}\right) + f\left(c_{\left(\frac{n}{2}+1\right)}\right) \text{ if } f\left(c_{\left(\frac{n}{2}\right)}\right) < f\left(c_{\left(\frac{n}{2}+2\right)}\right) \\ f\left(c_{\left(\frac{n}{2}+1\right)}\right) + f\left(c_{\left(\frac{n}{2}+2\right)}\right) \text{ if } f\left(c_{\left(\frac{n}{2}\right)}\right) > f\left(c_{\left(\frac{n}{2}+2\right)}\right) \end{array} \right\}$$

$$f(p_2) = f(p_1) + f(c_1)$$

$$f(p_i) = f(p_{(i-1)}) + f(p_{(i-2)}) \text{ for } 3 \leq i \leq m$$

$$f(v_1) = f(p_m) + f(p_{(m-1)}) \quad f(v_2) = f(v_1) + f(p_m)$$

$$f(v_i) = f(v_{(i-1)}) + f(v_{(i-2)}) \text{ for } 3 \leq i \leq n$$

$$f(x) = f(v_n) + f(v_1) \text{ and } f(y) = f(v_n) + f(v_{(n-1)})$$

Thus, G is sum graph with sum number 2 when  $n \geq 6$

Hence, G is sum graph with sum number 2 for all  $n \geq 3$ .

Illustration: Sum labeling for the graph  $[C_6, P_2]$  is given in figure 3.2




Figure 3.2

### References

[1] F Harary, *Graph Theory*, Addison Wesley, Reading, Massachusetts, 1972.

[2] J A Gallian, A dynamic survey of graph labeling, *The Electronics Journal of Combinatorics*, vol. 16, DS6, 2009.

[3] F Harary, *Sum Graphs and Difference Graphs*, Congress Numerantium, No.72, pp. 101-108, 1990

[4] P T Chung and S M Lee, On the super edge-graceful spiders of even orders, *J Combinatorial Mathematics and Combinatorial Computing*, vol. 64, pp. 3-17, 2008.

[5] S C Shee, Y S Ho, The cardinality of path-union of n copies of a graph, *Discrete Math*, vol. 151, pp. 221-229, 1996.

[6] G Sethuraman and P Selvaraju, Gracefulness of arbitrary super subdivisions of graphs, *Indian Journal of pure and applied Mathematics*, vol. 32, pp. 1059-1064, 2001.

[7] K M Kathiresan and S Amutha, Arbitrary supersubdivisions of stars are graceful, *Indian Journal of pure and applied Mathematics*, vol. 35, pp. 81-84, 2004.

[8] J Gerard Rozario, *Sum labeling for some cycle related graphs*, Paper presented in National Seminar (UGC Sponsored) on Recent Trends in Distance in Graphs on 12<sup>th</sup> and 13<sup>th</sup> March, at ANJAC, Sivakasi, 2012.

[9] R Ponraj, J V Xavier and R Kala, Some new families of Pair Sum graphs, *Journal of Mathematics and Computer Science*, vol. 2, pp. 747-758, 2012.