

## Sterol Composition of the Indian Green Lipped Mussel *Perna viridis*

Anuradha V\*, Vineetha P V†, Amrutha Manoj‡, Stephy P S§, Byju K\*\*, Shameem K††, Vasundhara G‡‡, N C Kumar§§ and S M Nair\*\*\*

### Abstract

Sterol composition of green lipped mussel *Perna viridis* was analysed using GCMS. Cholesterol was found to be the dominant sterol (54.162% of the total sterol content). Other sterols such as Cholesta-5,22-dien-3-ol (3 $\beta$ ), ergosta-5,22-dien-3-ol (3 $\beta$ ,22E, 24S), 26,26-dimethyl-5,24 (28)-Ergostadien-3 $\beta$ -ol, 26-nor-5cholesten- 3 $\beta$ -ol, stigmasterol and  $\gamma$  sitosterol were also detected. Presence of phytosterols like stigmasterol and  $\gamma$  sitosterol underlines high nutritional potential and food value of this bivalve mollusk.

**Keywords:** *Perna viridis*, Kerala coast, Sterols, Food value

---

\* Inter University Center for Development of Marine Biotechnology, Cochin University of Science and Technology, (IUCDMB, CUSAT), Cochin 16, India; radhavattoni@gmail.com

† (IUCDMB, CUSAT); viniithapv15@gmail.com

‡ (IUCDMB, CUSAT); amruthamanoj16@gmail.com

§ (IUCDMB, CUSAT); ps.stephy@gmail.com

\*\* (IUCDMB, CUSAT); byjuponnana@gmail.com

†† (IUCDMB, CUSAT); shameem2012@gmail.com

‡‡ (IUCDMB, CUSAT); drvasundharag@gmail.com

§§ (IUCDMB, CUSAT); chandramohan.kumar@gmail.com

\*\*\* (IUCDMB, CUSAT); unnimurali@yahoo.com

## 1. Introduction

The green mussel *Perna viridis*, commonly known as the Indian green mussel is a native of the IndoPacific region, primarily distributed along the Indian and south east Asian coasts. They generally inhabit marine intertidal, sub tidal and estuarine environments with high salinity. *Perna viridis* is a characteristic species of the fauna of mid littorial and sub littorial zones. The species is regarded as an important edible commodity. It is a remarkable species in terms of its ability to reach very high biomass level, to withstand environmental pollutants, to colonise artificial marine habitats and to invade new geographic territories. Extensive studies have been carried out on this particular species and it was proved to be an excellent source of carbohydrates, proteins, lipids and essential vitamins. Bivalves as sea foods are considered important next to fish and prawns, from nutritional point of view. They have been consumed for thousands of years. So the earlier investigations were limited to the biological, nutritional and medicinal aspects of this edible bivalve [1]. Studies on the protein content on selected organs of *Perna viridis* at Tranquebar coastal waters, Tamilnadu [2], work on the nutraceutical property of *Perna* [3], evaluation of the seasonal variation in nutritional values of *Perna viridis* by Wafar et al 1976[4] and the antibacterial activity of the extracts of green Mussel (*Perna viridis*) [5] are worth discussing in this context.

Despite the interest as a dietary source, little detailed information exists on the steroidal composition of Mussel. Even though attempts were made by Duoli , 2007 [6] to study the steroidal composition of *Perna viridis*, collected from China, the sterols were identified by comparison with mixtures of sterols from the NewZealand Green Lipped mussel, which had been previously identified by GC-MS [7]. This study endeavours to investigate the steroids present in the marine bivalve mollusc *Perna viridis* and here in we provide the first GCMS data of the sterol composition of previously unsampled *Perna viridis* from southern coast of India.

## 2. Experimental

The specimens were collected from Chellanam of Cochin during 2012 March. The live organisms after collection was thoroughly cleaned in seawater and stored in glass containers kept frozen until the start of the extraction work. The organism was macerated and further extracted with fresh methanol. The extracts were combined; filtered and excess alcohol was evaporated *invacuo* below 40°C. The concentrated crude methanolic extract was then evaporated and extracted with ethyl acetate, to acquire the lipophilic fraction. The extract was evaporated and the residues were purified by column chromatography. The residue of the ethyl acetate extract was column chromatographed over silica gel (60-120 mesh) and eluted with hexane with increasing concentration of ethylacetate, to get the semi pure fractions and analysed by GCMS.

GCMS analyses of the samples were done using PerkinElmer Clarus 680 GCMS equipped with head space sampler. The oven temperature was programmed from 35°C to 280°C at the rate of 10° per minute and is hold for 10 minutes. The column used for the analysis was a 30mx0.25mm i.d. capillary column coated with a film of dimethylpolysiloxane containing 5% diphenylpolysiloxane (Elite 5 MS Column) An aliquot of 0.5µl of the sample was injected into the GC-MS system in the split mode (split ratio 1 : 50). Helium was used as the carrier gas with a flow rate of 1mLmin<sup>-1</sup>. The injector and temperature was optimized at 280°C. The MS operating parameters were as follows: ionization energy -70 eV; ion source temperature-200°C, solvent delay 4 min and scan range 40-600 u. Identification of the components was based on direct comparison of mass spectral data with NIST library version 2.1.

## 3. Results and Discussion

Our investigation of *Perna viridis* provided us with seven steroids, cholesta-5,22-dien-3-ol(3β), cholesterol, ergosta-5,22-dien-3-ol (3 β,22E, 24S), 26,26-dimethyl-5,24 (28)-ergostadien-3β-ol, 26-nor-5cholesten- 3β-ol, stigmasterol and γ sitosterol. The total ion chromatogram and the percentage composition are given in Figure 1 and table 1 respectively. The structures of the sterols identified and the mass spectrum corresponding to each peak are

given in Figure 2 and 3. The steroids were found to present in the order cholesterol> ergosta-5,22-dien-3-ol (3  $\beta$ ,22E, 24S)> Cholesta-5,22-dien-3-ol (3 $\beta$ )> 26,26-dimethyl-5,24 (28)-Ergostadien-3 $\beta$ -ol = 26-nor-5cholesten- 3 $\beta$ -ol>  $\gamma$  sitosterol> stigmasterol.




Figure 1: Total Ion Chromatogram of the sterols identified

RT	Sterols Identified	% sterols/total sterol content
30.58	Cholesta-5,22-dien-3-ol (3 $\beta$ )	5.938
31.32	Cholesterol	54.162
31.92	ergosta-5,22-dien-3-ol (3 $\beta$ ,22E, 24S)	13.480
32.93	26,26-dimethyl-5,24 (28)-Ergostadien-3 $\beta$ -ol	7.036
33.08	26-nor-5cholesten- 3 $\beta$ -ol	8.434
33.59	Stigmasterol	2.668
34.88	$\gamma$ sitosterol	8.279

Table 1: Percentage composition of sterols




Figure 2: Structure of sterols identified




Figure 3: Mass fragmentation of the sterols identified from *Perna viridis*

Investigations on De-novo sterol synthesis in marine mollusks have assumed that cholesterol is the principal sterol formed from acetate and Mevalonate pathway [8, 9]. Composition of sterols other than cholesterol occurring in molluscan tissues may be affected by the dietary sources of sterols [10]. Also Khan and Goad (1983)[11] have mentioned three possible sources of sterols in mollusks 1. Denovo synthesis by usual  $\Delta^5$  sterol biosynthesis route, 2. Accumulation of dietary sterols, 3. Interconversion of dietary sterols to sterols in body of organisms. Compared with other marine invertebrates, marine bivalve mollusks are unique as they contain a wide range of

sterols, besides cholesterol[12]. Plant sterols such as stigmasterol and sitosterol was identified in species studied. It was established that the sterol profile of the lipids in the marine mollusks reflects the sterol profile of the food source (zooplankton, dinoflagellates and algae) [7].

#### 4. Conclusion

*Perna viridis* is a found to be a proven source for a variety of rare sterols. Previous studies have proved that, bivalves are low in steroids (50mg/g wet weight) when compared to the traditional protein sources such as beef (100mg/g wet weight) [12]. Also it was previously proved that non-cholesterol sterols derived from plants could reduce blood serum cholesterol in humans [13]. We could conclude that this bivalve under study, exhibited high conversion efficiency and hence high food value, with respect to the steroidal profile considered in this study. Also the accomplishment of the biological activity studies of these isolated steroids would certainly result in lead molecules for the development of promising drugs in future.

#### Acknowledgments

The financial support received from the government of Kerala to establish new centre, Inter University Centre for Development of Marine Biotechnology (IUCDMB) is gratefully acknowledged.

#### References

- [1] P. S. Kuriakose and K. K. Appukuttan, "Technology of mussels culture," *Bulletin Central Marine Fisheries Research Institute*, vol.48, pp.70-75, 1996.
- [2] D. Soumady and S. Asokan, "A Study on Protein Content in Selected Organs of *Perna Viridis* at Tranquebar Coastal waters, Tamilnadu," *World Journal of Zoology*, vol.6(4), pp.360-363, 2011
- [3] K. B Jena, T. G Jagtap and X. N Verlecar, "Antioxidative Potential of *Perna Viridis* and Its Protective Role Against ROS Lipid peroxide and Protein Carbonyl," *Current Trends in Biotechnology and Pharmacy*, vol.4 (4), pp. 862-870, October 2010.

- [4] M. V. M Wafer, S. Vijayaraghawan and L. Krishnakumar, "Seasonal Changes in the nutritive values of Green Mussel *Mytilus Viridis*," *Indian J. Marine Sci*, vol.5, pp.252-254, 1976.
- [5] Annamalai *et al*, "Antibacterial Activity of green Mussel (*Perna Viridis*) and Edible Oyster *Madorasensis*," *Research Journal of microbiology*, vol.2912), pp.978-982, 2007.
- [6] Duo Li *et al*, "Seasonal Variations of Lipid Content and Composition In *Perna Viridis*," *Lipids*, vol.42, pp.739-747, 2007.
- [7] K. Murphy, B. D Mooney, P. D Nichols and A.J Sinclair, "Lipid, Fatty Acid and Sterol Composition of Frozen-Dried New Zealand Green Lipped Mussel (*Perna Viridis*) and Tasmanian Blue Mussel (*Mytilus Edulis*)," *Lipids*, vol.37, pp.587-595, 2002.
- [8] D. R Idler and P. Wiseman, "Sterols of Crustacea" *Int. J. Biochem*, vol.2, pp. 91-98, 1971
- [9] S. Teshima, G. W. Patearnson "Sterols of Crustaceans, Molluscks and Fish in Physiology and Biochemistry of Sterols," *American Oil Chemists Society*, 1991.
- [10] T. Ando, A. Kanazawa, S. Teshima and H. Miyawaki, "Sterol Component of coral Reef Molluscks," *Marine Biology*, vol.50, pp.169-173, 1979.
- [11] A.S. Kahan and L.J Goad, "Sterol Constituents and Sterol Content of some Bivalve Molluscks," *Comp.Biochem.Physiology*, vol.76b, pp.569-573, 1983. doi:10.1016/0305-0491(83)90295-X
- [12] D. T. Gordon, "Sterols in Molluscks and Crustaceans of the Pacific Northwest," *J. Am. Oil Chem.Soc.*, Vol.59, pp.536-545, 1982.
- [13] V. Piironen, J. Toivo and A.M Lampi, "Natural Sources of Dietary Plant sterol," *Food Chem. Anal.*, Vol.13, pp.619-624, 2000. doi:10.1006/jfca.2000.0898